

Supremacy distribution in evolving networks

Janusz A. Hołyst,* Agata Fronczak,† and Piotr Fronczak
Faculty of Physics and Center of Excellence Complex Systems Research
Warsaw University of Technology
Koszykowa 75, PL-00-662 Warsaw, Poland
 (Dated: August 27, 2003)

We study a supremacy distribution in evolving Barabasi-Albert networks. The supremacy s_i of a node i is defined as a total number of all nodes that are younger than i and can be connected to it by a directed path. For a network with a characteristic parameter $m = 1, 2, 3, \dots$ the supremacy of an individual node increases with the network age as $t^{(1+m)/2}$ in an appropriate scaling region. It follows that there is a relation $s(k) \sim k^{m+1}$ between a node degree k and its supremacy s and the supremacy distribution $P(s)$ scales as $s^{-1-2/(1+m)}$. Analytic calculations basing on a continuum theory of supremacy evolution and on a corresponding rate equation have been confirmed by numerical simulations.

PACS numbers: 89.75.-k, 02.50.-r

I. INTRODUCTION

During the last few years there has been a large interest in modeling of networks [1, 2, 3, 4, 5] and several parameters describing the network structure have been considered. The examples are: degree distribution $P(k)$ [2, 6], mean path length [7, 8, 9, 10], betweenness centrality (load) [9, 11] or first and higher order clustering coefficients [12, 13, 14]. Universal scaling has been observed for some of these parameters in computer simulations and in real data describing such objects as the Internet, WWW, scientific collaboration networks or food webs [3, 4, 5]. Here we introduce a new parameter that can play an important role for description of a class of directed networks. We name the parameter a *supremacy* since it describes the number of nodes that are subordinated to a certain node. In the next Section we define our parameter and show its relevance for different problems of complex networks, Sec. III includes a continuum theory for the supremacy time evolution $s_i(t)$ and the supremacy probability distribution $P(s)$ in the Barabasi-Albert (BA) model with $m = 1$, in Sec. IV we find and solve a corresponding rate equation while in Sec. V a generalization of our problem for the BA model with $m > 1$ is presented.

II. THE MODEL

Let us consider the BA network with the characteristic parameter $m = 1$ [1, 2]. At the moment t_i a node i is created and it attaches to some older node in the network according to the preferential attachment rule (PAR). Then in the next time steps other nodes are created and are attached to the node i or to other nodes of




FIG. 1: Schematic illustration of the supremacy effects in tree-like BA network with $m = 1$. Numbers situated in the vicinity of the nodes represent their supremacies.

the network following PAR. As a result at the moment $t > t_i$ there is a subgraph in a form of tree $T(i, t)$ beginning in the node i and containing all nodes that are younger than the node i and that are connected to i by directed paths as in Fig. 1. If we assume that the node i represents a scientist who wrote an important paper or a politician who created an influential party [15, 16] we can consider all nodes belonging to the tree as his/her successors. If the tree $T(i, t)$ contains s_i nodes then the number s_i is the measure of the supremacy or the predominance of the node i at time t . The subgraph $T(i)$ can be also interpreted as a cluster of connected sites in the directed percolation problem [17, 18, 19, 20, 21, 22] and the supremacy of a node i is just the size of such a cluster starting from the site i . Since the evolution of the network is governed by PAR and all properties of the network are described by some probability distributions we are interested in the supremacy distribution $P(s)$ in the network.

*Electronic address: jholyst@if.pw.edu.pl

†Electronic address: agatka@if.pw.edu.pl




FIG. 2: Supremacy as a function the node degree. Solid lines represent analytical predictions of $s(k)$ given by Eq.5 in the case of $m = 1$ and Eq.13 in the case of $m = 2$.

III. CONTINUUM THEORY OF SUPREMACY EVOLUTION AND DISTRIBUTION FOR $m = 1$

To find the supremacy distribution $P(s)$ we follow the method that was introduced in [2] for calculation of degree distribution $P(k)$ in evolving networks. We start from determining the time dependence of $s_i(t)$ assuming that it is a continuous real variable. The supremacy of the node i increases in time because new nodes can be attached to any node of the tree $T(i, t)$. Let nodes belonging to the tree $T(i, t)$ possess degrees $k_i^{(1)}, k_i^{(2)}, \dots, k_i^{(s_i)}$.

Using the PAR we can write the following equation for changes of $s_i(t)$

$$\frac{\partial s_i(t)}{\partial t} = \sum_{l=1}^{s_i} \frac{k_i^{(l)}}{2t} = \frac{K(i)}{2t}, \quad (1)$$

where $K(i) = \sum_{l=1}^{s_i} k_i^{(l)}$ and we used the fact that at the moment t the sum of all nodes degrees in the whole network equals to $2t$. On the other hand taking into account the tree structure of the considered subgraph we can write the supremacy s_i as

$$s_i = 1 + \sum_{l=1}^{s_i} (k_i^{(l)} - 1) = 1 + K(i) - s_i, \quad (2)$$

thus $K(i) = 2s_i - 1$ and we have a simple equation

$$\frac{\partial s_i(t)}{\partial t} = \frac{2s_i - 1}{2t}, \quad (3)$$

with the solution

$$s_i(t) = \frac{1}{2} \left(\frac{t}{t_i} + 1 \right), \quad (4)$$

where we took into account the initial condition $s_i(t = t_i) = 1$. The solution (4) means that the node supremacy




FIG. 3: Supremacy distribution in BA model. Solid lines represent analytical predictions of $P(s)$ given by Eq.7 in the case of $m = 1$ and Eq.14 in the case of $m = 3, 5$.

increases linearly in time comparing to the square root dependence of the node degree [2], i.e. $k_i(t) = \sqrt{\frac{t}{t_i}}$. Combining the last two results we get a simple relation between the node supremacy and the node degree

$$s(k) = \frac{1}{2} [k^2 + 1]. \quad (5)$$

In the region $k \leq 100$ this formula fits well to numerical simulations presented in Fig.2 while for larger k differences between the analytic theory and the numerical simulations are observed.

The probability density $P(s)$ for the supremacy distribution in the network follows from the relation

$$P(s_i) ds_i = \tilde{P}(t_i) dt_i, \quad (6)$$

where $\tilde{P}(t_i) = 1/t$ is the distribution of nodes attachment times t_i for a network of age t . After a simple algebra we get

$$P(s_i) = \frac{1}{t} \left| \frac{\partial s_i}{\partial t_i} \right|^{-1} = \frac{2}{(2s_i - 1)^2}. \quad (7)$$

One can see that the supremacy distribution is a time independent function. Fig.3 shows the comparison of the last equation to numerical data. Let us stress that for $s \gg 1$ the supremacy distribution scales as $P(s) \sim s^{-2}$ while the degree distribution for BA model [1, 2] scales as $P(k) \sim k^{-3}$.

IV. RATE-EQUATION FOR SUPREMACY DISTRIBUTION FOR $m = 1$

Now we show how to get the supremacy distribution using the rate-equation approach that was introduced by Krapivsky, Redner and Leyvraz [6] to study networks degree distribution $P(k)$. Let $N(s, t)$ is the number of

nodes possessing the supremacy s at time t . The rate equation for $N(s, t)$ is

$$\frac{\partial N(s, t)}{\partial t} = \frac{[2(s-1) - 1]N(s-1, t) - (2s-1)N(s, t)}{2t} + \delta_{s,1}. \quad (8)$$

The first term on the right-hand side of Eq. (8) corresponds to creation of a new node with the supremacy s . The process is proportional to the number of nodes with the supremacy $s-1$ and the corresponding transition probability that follows from the PAR and Eq. (2). The second term corresponds to creation of a node with a supremacy $s+1$, i.e. to destruction of a node with a supremacy s while the last term describes creation of a node with a supremacy $s=1$. Writing $N(s, t) = P(s)N_0$ where $N_0 = t$ corresponds to the total number of nodes at time t and $P(s)$ is the probability of a node with the supremacy value s we get the recursive equation

$$P(s) = \frac{2s-3}{2s+1}P(s-1) \quad \text{for } s \geq 2, \quad (9)$$

where $P(1) = 2/3$. The solution of Eq. (9) is

$$P(s) = \frac{2}{(2s-1)(2s+1)}. \quad (10)$$

Note that for $s \gg 1$ the solution (10) coincides with the solution (7) that has been received in the limit of the continuum theory.

V. SCALING OF SUPREMACY DISTRIBUTION FOR $m > 1$

The peculiar feature of the BA model is the independence of the scaling exponent characterizing the degree distribution $P(k) \sim k^{-3}$ from the model parameter m describing the number of links that are created by every new node. Below we show that the scaling exponent of supremacy distribution depends on the parameter m . If we neglect all loops existing in the BA network with the characteristic parameter $m > 1$ then we can easily repeat our considerations from Sec.III and IV. Instead of Eq. (2) we get

$$s_i = 1 + \sum_{l=1}^{s_i} (k_i^{(l)} - m) = K(i) + 1 - ms_i, \quad (11)$$

and time evolution of the supremacy is described by

$$s_i(t) = \frac{m}{m+1} \left(\frac{t}{t_i} \right)^{\frac{m+1}{2}} + \frac{1}{m+1}, \quad (12)$$

thus the relation between the degree and the supremacy is

$$s(k) = \frac{m}{m+1} \left(\frac{k}{m} \right)^{m+1} + \frac{1}{m+1}. \quad (13)$$




FIG. 4: Schematic illustration of supremacy effects in BA network with $m = 2$. Solid arrows represent connections within the supremacy area / cluster of the top vertex, whereas dashed arrows express connections pointing outside the cluster. Note that there is a single loop in the cluster.

It follows that for dense networks with $m \gg 1$ the supremacy $s_i(t)$ increases in time much faster than the node degree $k_i(t)$. Fig. 2 shows a comparison of the result (13) to numerical data for $m = 2$. One can see that the predicted scaling of $s(k)$ breaks down completely for large values of k where the plot $s(k)$ saturate. The reason is the presence of loops that for $m > 1$ appear in the network and that have been neglected in our approach. If $m > 1$ the result (13) is valid mainly for vertices with a small degree k_i (and a small supremacy s_i) since loops are sparse in small clusters starting from such nodes. The saturation effect does not appear for the BA model with the parameter $m = 1$ where loops are absent.

Taking into account (12) we get the supremacy distribution in the form

$$P(s_i) = \frac{1}{t} \left| \frac{\partial s_i}{\partial t_i} \right|^{-1} = \frac{2}{m} \left[\frac{(m+1)s_i - 1}{m} \right]^{-\frac{m+3}{m+1}}. \quad (14)$$

We see that the scaling exponent for the supremacy distribution equals to $\delta = -1 - 2/(1+m)$ and in contrast to the scaling exponent of degree distribution it depends on the parameter m . The result (14) is in a good agreement with numerical simulation for BA networks, see Fig. 3.

The rate-equation for $m > 1$ is similar to Eq. (8), i.e.

$$\frac{\partial N(s, t)}{\partial t} = \frac{[(1+m)(s-1) - 1]N(s-1, t)}{2t} - \frac{[(1+m)s - 1]N(s, t)}{2t} + \delta_{s,1} \quad (15)$$

The resulting solution for the probability $P(s)$ can be written as the following product

$$P(s) = \frac{2}{m+2} \prod_{i=2}^s \frac{[(i-1)(m+1) - 1]}{[i(m+1) + 1]} \quad (16)$$

for $s > 1$ where $P(1) = 2/(m + 2)$. For dense networks $m \gg 1$ the solution (16) can be approximately written as

$$P(s) \simeq \frac{2}{ms} \quad (17)$$

what coincides with (14).

VI. CONCLUSIONS

In conclusion, we introduced a universal parameter (a supremacy) that describes vertices in directed networks. The parameter equals to the size of a cluster starting from the site in a directed percolation model. We have shown that for Barabasi-Albert model there is a relation between the supremacy and the vertex degree. It

follows that there are universal scaling laws describing the time evolution of the supremacy and corresponding supremacy distributions in BA models. On the contrary to the scaling results for nodes degree the corresponding scaling exponents of supremacy depend on the characteristic model parameter m . Numerical simulations are in good agreement with analytical estimations for node with a small and medium supremacy especially for the case $m = 1$ where no loops are present in the system.

VII. ACKNOWLEDGMENTS

The paper was supported by the special program of the Warsaw University of Technology *Dynamics of Complex Systems*.

-
- [1] A.L. Barabási and R. Albert, *Science* **286**, 509 (1999).
 - [2] A.L. Barabási, R. Albert and H. Jeong, *Physica A* **272**, 173 (1999).
 - [3] *Handbook of Graphs and Networks*, edited by S. Bornholdt and H.G. Schuster (Wiley-Vch, Weinheim, 2002).
 - [4] S.N. Dorogovtsev and J.F.F. Mendes, *Evolution of Networks*, Oxford Univ.Press (2003).
 - [5] R. Albert and A.L. Barabasi, *Review of Modern Physics* **74**, 47 (2002).
 - [6] P.L. Krapivsky, S. Redner and F. Leyvraz, *Phys. Rev. Lett.* **85**, 4629 (2000).
 - [7] M.E.J.Newman, C.Moore and D.J.Watts, *Phys. Rev. Lett.* **84** 3201 (2000).
 - [8] M.E.J.Newman, S.H.Strogatz and D.J.Watts, *Phys. Rev. E* **64**, 026118 (2001).
 - [9] G.Szabó, M.Alava and J.Kertész, *Phys. Rev. E* **66** 026101 (2002).
 - [10] A. Fronczak, P. Fronczak and J.A. Hołyst, cond-mat/0212230.
 - [11] K.-I. Goh, B. Kahng, and D. Kim, *Phys. Rev. Lett.* **87**, 278701 (2001).
 - [12] D.J.Watts and S.H.Strogatz, *Nature* **393**, 440 (1998).
 - [13] A. Fronczak, P. Fronczak and J.A. Hołyst, cond-mat/0306255, *Phys. Rev. E* (in print).
 - [14] A. Fronczak, J.A. Hołyst, M. Jedynak and J. Sienkiewicz, *Physica A* **316**, 688 (2002).
 - [15] K. Kacperski and J.A. Hołyst, *J. Stat. Phys.* **84**, 169 (1996).
 - [16] K. Kacperski and J.A. Hołyst, *Physica A* **269**, 511 (1999); *Physica A* **287**, 631 (2000).
 - [17] P. Grassberger, *Journ. Phys. A - Math. General* **22**, 3673 (1989); *Journ. Stat. Phys.* **79**, 13 (1995).
 - [18] A. Bunde and S. Havlin, in *Fractals and Disordered Systems*, edited by A. Bunde and S. Havlin (Springer, Heidelberg, 1996), 2nd ed.
 - [19] R. Cohen, S. Havlin, and D. ben-Avraham, in *Handbook of Graphs and networks*, edited by S. Bornholdt and H.G. Schuster (Wiley-Vch, Weinheim, 2002).
 - [20] A. Aleksiejuk and J.A. Hołyst, *Physica A* **299**, 198 (2001).
 - [21] M. E. J. Newman, S. H. Strogatz and D. J. Watts, *Phys. Rev. E* **64**, 026118 (2001).
 - [22] S. N. Dorogovtsev, J. F. F. Mendes and A. N. Samukhin, *Phys. Rev. E* **64**, 025101(R) (2001).